

APPLICATION
FOR
UNITED STATES LETTERS PATENT

TITLE: TOXIN-PHAGE BACTERIOCIDE ANTIBIOTIC AND USES
THEREOF

APPLICANT: DIANE L. SCHAAK

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL932 079 390US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

December 18, 2001
Date of Deposit

[Signature]
Signature

Lisa G. Gray
Typed or Printed Name of Person Signing Certificate

Toxin-Phage Bacteriocide Antibiotic and Uses Thereof

TECHNICAL FIELD

This invention relates to compositions and methods for killing bacteria.

BACKGROUND

Throughout recorded history virulent bacterial infections have been a bane to mankind. Until recently, it was assumed that drug antibiotics had largely eradicated virulent bacteria. It is now apparent, however, that bacteria have circumvented the effects of single-point targeted drug antibiotics. Consequently, there is a need to develop new anti-bacterial agents that can be used to supplement or replace conventional drug antibiotics.

Like animal cells, bacterial cells are subject to infectious agents that are present in their environment. Viruses known as bacteriophage, or phage, specifically infect bacterial cells. Bacteriophage are the natural enemies of bacteria and, over the course of evolution, have developed proteins which enable them to infect a bacterial host cell, replicate their genetic material, usurp host metabolism, and ultimately kill their bacterial host cell.

Research into the use of bacteriophage as therapeutic agents for treatment of bacterial infection began sometime in the late 19th century, predating the development of conventional drug antibiotics. By 1920, Edward Twort and Felix d'Herelle, two noted pioneers in bacteriophage research, were isolating bacteriophage from several bacterial species and using them as anti-bacterial agents. During the early 1940's, however, antibiotics were introduced to the world as a broad range treatment for bacterial infections, and bacteriophage therapy research went into decline.

Early clinical studies of phage therapy were plagued with poor experimental design, with few controls and little documentation, variable success due to the indiscriminate use of phage to treat a broad range of bacterial infections, and the use of procedures that introduced bacterial toxins into patients and loss of effectiveness of the isolated phage.

The lack of knowledge and scientific expertise needed to understand bacteriophage and their interaction with bacteria also hindered efforts to improve phage therapy. For example, differences between the biological interaction of bacteriophage strains with their species-specific bacterial host *in vitro* as compared to *in vivo* have posed considerable difficulty. Although bacteriophage can be selected for their lytic virulence (immediately

replicating and then inducing bacterial host cell lysis following infection) *in vitro*, such selection does not guarantee against the conversion of a seemingly lytic phage to a temperate phage (entering into a state of lysogeny *via* integration of the bacteriophage genome into the bacterial genome followed by a quiescent period during which lytic proteins are not expressed) *in vivo*. These conversions result in lysogenic bacteria that are resistant to further bacteriophage infection, thus reducing the effectiveness of phage therapy.

Since the early 1940's drug antibiotics have become the choice for treating virulent bacterial infections. Several problems associated with this approach are now becoming evident. The misuse and overuse of drug antibiotics has contributed to the rise of antibiotic resistant bacterial strains. Moreover, since drug antibiotics are non-specific with respect to the types of bacteria that they effect, the bacterial flora that naturally occur within the body are killed along with the disease-causing bacterial pathogen. At least 200 identified bacterial species normally inhabit the human body, and many of the these species synthesize and excrete vitamins vital for human health, promote the development of certain tissues, e.g., lymphatic tissue, e.g., Peyer's patches, and stimulate the production of cross-reactive "natural" antibodies that react with pathogenic bacteria. Moreover, natural bacterial flora greatly inhibit colonization by non-indigenous bacteria through normal niche colonization or by producing substances and bacteriocins that can inhibit and kill foreign bacteria. Conventional broad spectrum antibiotics risk killing the non-pathogenic bacteria that are responsible for these beneficial effects.

Bacterial drug resistance was evident at the onset of drug antibiotic therapy, and drug resistant virulent strains of both gram-negative bacteria (including pathogenic strains of *Escheria coli*) and gram-positive bacteria (including pathogenic strains of *Staphylococcus* and *Streptococcus*) have become increasingly resistant to drug antibiotics. This increased resistance arises primarily from selection for virulent-resistance strains by the presence of drug antibiotics, resulting in the lateral transfer of resistance genes between different strains and species of bacteria. Epidemic outbreaks have been attributed to a single clone of a benign or virulent progenitor, as well as spontaneous multi-clonal populations within a community setting when drug antibiotic usage is increased. Although decreased usage of antibiotics may improve the odds of generating a population of virulent bacteria that are less resistance towards antibiotics, much contradictory evidence is beginning to surface. For

example, a study in Finland found that the incidence of *Streptococcus pyogenes* resistance to macrolide decreased after macrolide treatment was reduced in favor of treatment with erythromycin. However, a follow-up study reported a subsequent 17% increase in *Streptococcus pyogenes* resistance to erythromycin. Another growing concern is the increasing number of multi-resistant bacteria. In 1968 approximately 12,500 people in Guatemala died from an epidemic of Shigella, caused by a bacterial strain that contained a plasmid encoding genes resistant to four different antibiotics (Davies (1996) *Nature* 383:219). Population genetics studies of virulent bacteria causing disease outbreaks or increases in frequency and virulence have shown that the distinct clones responsible for the acute outbreaks are often characterized by unique combinations of virulence genes or alleles of those genes.

Increasing drug antibiotic resistance has resulted in increased dosage levels and duration of antibiotic treatment. These practices are associated with hypersensitivity and serious side effects in a growing number of patients (see Cunha (2001) *Med Clin North Am* 85:149; Kirjavainen and Gibson (1999) *Ann Med* 31:288; Lee et al. (2000) *Arch Intern Med* 160:2819; and Martinez et al. (1999) *Medicine* 78:361). The increasing hypersensitivity and side effects are not being seriously addressed and have so far been clinically under-evaluated (Demoly et al. (2000) *Bull Acad Natl Med* 184:761; and Gruchalla (2000) *Allergy Asthma Proc* 21:39). As an example of one serious side effect that is becoming increasingly prevalent, especially in children, the use of antibiotics has been shown to be positively associated with the development of asthma and atopy. The mechanisms underlying these associations remain largely unknown (von Hertzen (2000) *Ann Med* 32:397).

Drug antibiotics and their effects are not isolated to individuals under the supervision of a doctor's care, but are a communal health issue. Molecular population studies have identified healthy humans that are VRE (vancomycin-resistant enterococci) carriers. An increase in VRE strains in healthy farm animals is associated with the increased use of the antibiotic avoparcin. There is currently a tentative link between the consumption of farm animals and VRE transference to people (Bates (1998) *J Hosp Infect* 27:89). Data on antibiotic resistance profiles of several food born pathogens provides ample evidence that antibiotic resistance traits have entered the microflora of farm animals and the food supply produced from them (Teuber (1999) *Cell Mol Life Sci* 56:755).

SUMMARY

The present invention is based, at least in part, on the development of intracellular peptide toxins and peptide-like toxins that are toxic to a cell when inside the cell, but relatively non-toxic to the cell when outside the cell. Such peptide toxins and peptide-like toxins are useful in the production of a recombinant bacteriophage that effectively function as a bacteriocide (i.e., a toxin-phage bacteriocide) that can provide a viable alternative to conventional drug antibiotics. The toxin-phage bacteriocide (TPB) include bacteriophage that have been genetically engineered to encode a peptide toxin that can be expressed within the bacterial host cell. Within the bacterial host cell, the peptide toxin is active and functions to kill the bacterial host cell. Importantly, the toxin-phage bacteriocide of the invention retains its activity as a bacteriophage, and is therefore capable of completing the lytic phase of its lifecycle. Completion of the lytic phase of its life-cycle results in both the production of additional toxin-phage bacteriocide and host cell lysis.

Accordingly, in one aspect, the invention features a method of producing a toxin-phage bacteriocide. The method includes: (a) identifying a bacteriophage that is capable of infecting a bacterial cell of interest; (b) preparing a recombinant bacteriophage genome via the introduction of a nucleic acid sequence that encodes an intracellular peptide toxin into the genome of the bacteriophage, wherein the nucleic acid sequence that encodes the peptide toxin is operatively linked to a promoter that is active within the bacterial cell of interest; and (c) allowing the formation of a toxin-phage bacteriocide particle that contains the recombinant bacteriophage genome.

In preferred embodiments, the nucleic acid sequence that encodes an intracellular peptide toxin includes the nucleic acid of SEQ ID NO:1, which encodes the TPB peptide toxin A amino acid sequence (SEQ ID NO:2). In other embodiments, the nucleic acid sequence that encodes an intracellular peptide toxin encodes a peptide toxin other than the TPB peptide toxin A, e.g., a peptide toxin that is a variant of the amino acid sequence of TPB peptide toxin A, or a peptide toxin that functions analogously to the TPB peptide toxin A. In some embodiments, a variant of the TPB peptide toxin A includes at least one mutation, e.g., an insertion, deletion, or point mutation. In preferred embodiments, the mutation is located at one or more of amino acids 16, 17, 18, 19, 20, 21, and 22 of SEQ ID NO:2. In other

5

[REDACTED]

10

15

5

genome. In related embodiments, homologous recombination is carried out *in vitro*. In other related embodiments, homologous recombination is carried out *in vivo*. In other embodiments, the recombinant bacteriophage genome is packaged into bacteriophage particles *in vitro* or *in vivo*, thereby resulting in the production of toxin-phage bacteriocide particles.

In a related aspect, the invention features compositions that include at least one toxin-phage bacteriocide. In preferred embodiments, the toxin phage bacteriocide includes a nucleic acid sequence encoding an intracellular peptide toxin. In particularly preferred embodiments, the toxin phage bacteriocide includes a nucleic acid sequence encoding the TPB peptide toxin A (SEQ ID NO:2). In other embodiments, the toxin phage bacteriocide includes a nucleic acid sequence encoding TPB peptide toxin A variants.

In preferred embodiments, the compositions include a single strain or multiple variant strains of toxin-phage bacteriocide that has been substantially purified away from the bacterial host cells used to produce or amplify the toxin-phage bacteriocide. In other preferred embodiments, the compositions include a toxin-phage bacteriocide that has been substantially purified away from the bacterial host cell medium in which the bacterial host cells were grown during the production or amplification of the toxin-phage bacteriocide. In other embodiments, the compositions include a toxin-phage bacteriocide that has been partially purified from the bacterial host cells and bacterial host cell medium used to produce or amplify the toxin-phage bacteriocide.

In another aspect, the invention features a method of using a toxin-phage bacteriocide to kill a bacterial cell. The method involves contacting bacterial cells (e.g., bacterial cells that include one or more strains or species of bacteria) with a toxin-phage bacteriocide, such that at least one toxin-phage is able to bind to and infect at least one bacterial cell, and then allowing the toxin-phage that have infected bacterial cells to kill the bacterial cells. In preferred embodiments, the toxin-phage binds to and infects bacterial cells that are of a selected type. In other preferred embodiments, the toxin-phage does not bind to or infect bacterial cells that are not of the selected type. The contacting can occur within a patient, e.g., a human or animal patient, or *in vitro*. *In vitro* studies using the gram negative *Escheria coli* and gram positive *Bacillus subtilis* have found a 100% non-infectivity in the presence of a foreign toxin-phage.

In some embodiments, an infected bacterial cell is killed as a result of the toxin-phage entering into the lytic phase of its life-cycle, such that the bacterial cell is killed by lysis. In other embodiments, the infected bacterial cell is killed as a result of the expression of the toxic peptide encoded by the nucleic acid molecule that was introduced into the genome of the toxin-phage. In still other embodiments, the bacterial cell is killed by a combination of the toxin-phage entering into the lytic phase of its life-cycle and the expression of the toxic peptide encoded by the nucleic acid molecule that was introduced into the bacterial cell by the toxin-phage. In other embodiments, a bacterial cell that is killed is either a gram-negative or a gram-positive bacterial cell.

In another aspect, the invention features a pharmaceutical composition that includes at least one toxin-phage bacteriocide and at least one pharmaceutically acceptable carrier. In preferred embodiments, the pharmaceutical composition can be used *in vivo*, e.g., the pharmaceutical composition can be administered, e.g., by parenteral injection or orally, to a subject, to treat a bacterial infection present in the subject. In other embodiments, the pharmaceutical composition can be used topically to treat a bacterial infection present in or on a subject.

In another aspect, the invention features a method of using a toxin-phage bacteriocide to treat a bacterial infection present in or on a subject. In some embodiments, the subject is a farm animal, e.g., a chicken, pig, goat, sheep, cow, or horse. In other embodiments the subject is a plant, e.g., an agricultural product or orchard tree. In other embodiments, the subject is a pet, e.g., a fish, bird, cat, or dog. In still other embodiments, the subject is a mammal, a primate, or a human. In preferred embodiments, the toxin-phage bacteriocide kills the bacteria that are the cause of the infection. In other embodiments, the toxin-phage bacteriocide slows or brings to a halt the spread of the bacterial infection. In preferred embodiments, the toxin-phage bacteriocide helps eliminate the bacterial infection. In other preferred embodiments, the toxin-phage bacteriocide does not kill the bacterial cells that are not the cause of the infection, e.g., bacterial cells that are normally present in the subject or are beneficial to the subject. In other embodiments, the infection constitutes a localized disease, e.g., a disease of the skin, nervous system, cardiovascular system, respiratory system, digestive system, and urinary and reproductive systems.

In another aspect, the invention features a method of using a toxin-phage bacteriocide to prophylactically treat a potential bacterial infection in a subject. In some embodiments, the subject is a farm animal, e.g., a chicken, pig, goat, sheep, cow, or horse. In other embodiments the subject is a plant, e.g., an agricultural product or orchard tree. In other
5 embodiments, the subject is a pet, e.g., a fish, bird, cat, or dog. In still other embodiments, the subject is a mammal, a primate, or a human. In preferred embodiments, the toxin-phage bacteriocide kills the bacteria that are the potential cause of infection. In other embodiments, the toxin-phage slows or brings to a halt the growth of the bacteria that are the potential cause of infection. In other preferred embodiments, the toxin-phage bacteriocide does not kill
10 bacterial cells that are not the potential cause of infection, e.g., bacterial cells that are normally present in the subject or are beneficial to the subject. In other embodiments, the potential bacterial infection can result in acne, e.g., skin acne in a human. In other
15 embodiments, the subject has an injury, e.g., a cut that breaks the outer dermal layer of the skin, an animal bite, a dermal burn, or a surgical wound or incision, or a surgically inserted device, e.g., a catheter, that is highly susceptible to bacterial infection. In still other
embodiments, the potential bacterial infection can involve exposure to biological weapons, e.g., anthrax, plague, or tularemia.

In another aspect, the invention features a method of treating an aqueous solution with a toxin-phage bacteriocide such that bacteria present in the solution are killed. In one
20 embodiment, the resulting aqueous solution is partially sterilized and can subsequently be consumed by an animal, e.g., a farm animal, pet, mammal, primate, or human. Treatment of the aqueous solution will reduce the chance of bacterial infection resulting from consumption of the solution. In another embodiment, the aqueous solution is a solution that is subject to bacterial contamination, e.g., the water in a fish tank or wastewater, e.g., sewage.

25 In another aspect, the invention features a method of treating a surface with one or more toxin-phage bacteriocides such that bacteria attached to the surface are killed or their growth is inhibited. In one embodiment, the surface is part of a device, e.g., a device that is used in medicine (e.g., surgical instruments), agriculture, industrial processes, or water and wastewater treatment. In another embodiment, the surface is covered with a biofilm. In
30 other embodiments, the surface is treated regularly with a toxin-phage bacteriocide such that the formation of a biofilm is prevented or slowed.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

TPB can be designed to be specific for any selected strain of bacteria, thus desirable bacteria can be spared. Bacteriophage specific for a single bacterial host in nature have been found to remain within the host for as long as the bacterial host specific for that phage is present. Weber-Dabrowska, et al. (1987), *Arch Immunol Ther Exp* (Warsz) 35(5):563-8, tested for absorption of orally administered anti-staphylococcal and anti-pseudomonas phage in both urine and serum samples of patients with suppurative bacterial infections. No phage was present in any of the 56 patients prior to phage therapy. By day 10, 84% of the serum samples and 35% of urine samples contained phage, indicating bioavailability. The healthy control group exhibited a phage titer drop 100-fold between days 0-5. A comprehensive review of phage therapy (Alisky et al. (1998), *J of Infection* 36:5) concluded that all studies with both human and animals showed no measurable antiphage antibodies generated.

Without being bound by any particular theory, it appears that the TPB peptide toxin A, produced by a TPB of the invention, becomes introduced into internally available membranes of the cell. This has been observed to occur in both bacterial and yeast cells. *In vitro* studies using a lipid bilayer membrane model suggest that the toxin peptide permeabilizes membranes. Significantly, the TPB peptide toxin A does not appear to harm either bacterial cells or eukaryotic cells when applied externally, e.g., when introduced in a culture of growing cells.

The TPB peptide toxin A of the invention has also been found to be toxic to eukaryotic cell when presented internally. Thus, intracellular peptide toxins can be used to selectively target undesirable eukaryotic cells, e.g., cancer cells or virally infected cells, by selectively delivering the peptide toxins to the interior of the undesirable cells. Thus, the peptide toxins can be targeted to such cells in various ways, e.g., through receptor mediated targeting.

This invention is further illustrated by the following examples that should not be construed as limiting.

Example 1: Production of a Toxin Gene Master Stock

A nucleic acid molecule encoding the TPB peptide toxin A can be prepared synthetically. The molecule has the sequence: ATG GAT TGG CTG AAA GCT CGG GTT GAA CAG GAA CTG CAG GCT CTG GAA GCA CGT GGT ACC GAT TCC AAC GCT GAG CTG CGG GCT ATG GAA GCT AAA CTT AAG GCT GAA ATC CAG AAG (SEQ

ID NO:1). The nucleic acid molecule encodes a 39 amino acid peptide having the sequence: MDWLKARVEQELQALEARGTDSNAELRAMEAKLKAEIQK (**SEQ ID NO:2**).

The TPB peptide toxin A encoding nucleic acid molecule (**SEQ ID NO:1**) was inserted into pET19b plasmid (Novagen, Inc.; Madison, WI). The expression vector BL21-Gold(DE3)plysS (Stratagene, Inc.; La Jolla CA) was used for expression of the TPB peptide toxin A for *in vitro* studies. A TPB peptide toxin A encoding gene can be prepared by PCR amplifying a TPB peptide toxin A encoding nucleic acid molecule out of the pET19b plasmid, as discussed below, or by PCR amplification from a synthetically prepared nucleic acid molecule.

The top strand 5' oligonucleotide (**SEQ ID NO:3**) used for PCR amplification of the TPB peptide toxin A encoding gene included: an multiple cloning site (MCS), a promoter sequence that is functionally active in both gram-negative and gram-positive bacterial hosts, and a sequence homologous to the 5' start region of the toxin gene sequence. It had the following sequence:

GCGTCCGGCGTAGAGGATCCAAGCTTTAATTAAATTTTATTTGACAAAAATGGG
CTCGTGTTGTACAAATGTATGGATTGGCTGAAAGCTCGGGTTGAACAGG (**SEQ ID NO:3**). The first underlined portion is the MCS sequence. Restriction endonucleases that are capable of cutting within this MCS sequence are shown in Table I.

Table I

Enzyme	No.	Position	Sequence
AclWI	1	19	ggatc
AluI	1	23	ag/ct
AlwI	1	19	ggatc
BamHI	1	15	g/gatcc
BsiSI	1	5	c/cgg
Bsp143I	1	15	/gatc
BstI	1	15	g/gatcc
BstX2I	1	15	r/gatcy
BstYI	1	15	r/gatcy
CviJI	1	23	rg/cy
DpnI	1	17	ga/tc
DpnII	1	15	/gatc
HapII	1	5	c/cgg
HgaI	1	5	gacgc
HindIII	1	21	a/agctt
HpaII	1	5	c/cgg

	Kzo9I	1	15	/gatc
	MboI	1	15	/gatc
	MflI	1	15	r/gatcy
	MnlI	1	16	cctc
5	MseI	1	26	t/taa
	MspI	1	5	c/cgg
	NdeII	1	15	/gatc
	NlaIV	1	17	ggn/ncc
	PspN4I	1	17	ggn/ncc
10	Sau3AI	1	15	/gatc
	Sse9I	1	27	/aatt
	TruII	1	26	t/taa
	Tru9I	1	26	t/taa
	Tsp509I	1	27	/aatt
15	TspEI	1	27	/aatt
	XhoII	1	15	r/gatcy

The central portion of the top strand 5' oligonucleotide sequence (**SEQ ID NO:3**), which is not underlined, constitutes the VegI/II promoter sequence. The VegI/II promoter sequence has been shown by Pescheke et al. (1985), *J Mol Biol* 186:547, to be active in both gram-negative and gram-positive bacterial cells. The second underlined portion of the top strand 5' oligonucleotide sequence corresponds to the 5' end of the TPB peptide toxin A gene sequence (**SEQ ID NO:1**). This sequence is capable of annealing to the bottom strand of the pET19b plasmid, e.g., in a PCR reaction.

The bottom strand 3' oligonucleotide (**SEQ ID NO:4**) used for PCR amplification of the TPB peptide toxin A encoding gene included a MCS site and a sequence complementary to the 3' end of the toxin gene sequence. The terminator region present in the pET19b vector was not amplified so that the functional properties of the toxin peptide could be disrupted, rendering the gene product less toxic to the master stock host cell. The bottom strand 3' oligonucleotide used for PCR amplification had the sequence:

CCATCGATGGCCGCTCGAGCTATTATTTCTGGATTTCAG (**SEQ ID NO:4**). The underlined portion of **SEQ ID NO:4** constitutes the multiple cloning sites (MCS) sequence. Restriction endonucleases that are capable of cutting within this MCS sequence are shown in Table II.

Table II

20
26
30

35

40

GGCGTATCACGAGGCCC (SEQ ID NO:5); and GTGGCGCCGGTGATGCCGG (SEQ ID NO:6). SEQ ID NO:5 was used to sequence the PCR product from the 5' direction, while SEQ ID NO:6 was used to sequence the PCR product from the 3' direction.

The purified PCR product was cut with the restriction endonucleases ClaI and BamHI and ligated into a pBR322 plasmid (ATCC 37017, 31344) that had been cut with the same enzymes. Insertion of the PCR product containing the TPB peptide toxin A gene PCR product into the pBR322 plasmid disrupted the tetR gene, negating tetracycline resistance. This disruption, in turn, allowed for a positive gene incorporation selection tool. Once a positive clone was identified, the region of the plasmid containing the TPB peptide toxin A gene PCR product was analyzed using restriction digests, and then sequenced.

The resulting plasmid was transformed into competent HB101 (MAX Efficiency HB101 Competent Cells, Cat. No. 18296-012, Life Technologies), and a positive clone was chosen using ampicillin resistance as a selection criteria. A single colony clone was selected and cultured to exponential growth phase (LB, 37°C, 250 rpms), mixed with sterile glycerol (80:20 ratio) and stored in a -76°C freezer.

Example 2: Selection of Toxin Gene Integration Sites

Both a gram-negative and a gram-positive bacterial species with their complimentary bacteriophages were chosen to illustrate the effectiveness of TPB peptide toxin A.

Escherichia coli (c600, ATCC Accession No. 23724) was chosen as an example of a gram-negative bacterial species that could be tested for the effects of a toxin-phage bacteriocide. There are many bacteriophage that are known to infect *E. coli*, one of which is lambda phage (ATCC Accession No. 23724-B2). The sequence of the lambda phage genome is described in Sanger et al. (1992) *J Mol Biol* 162:729, the contents of which are incorporated herein by reference. The integration site for the TPB peptide toxin gene into the lambda phage genome was chosen to be between nucleotides 46,468 and 46,469. The nucleotide sequences of the regions immediately surrounding the chosen integration site are as follows: TTGCCCATATCGATGGGCAACTCATGCAATTATTGTGAG (SEQ ID NO:7); and CAATACACACGCGCTTCCAGCGGAGTATAAATGCCTAAAGTA (SEQ ID NO:8). SEQ ID NO:7 corresponds to the nucleotide sequence that is 5' to the integration site, about nucleotides 46,430 – 46,468 of the lambda phage genome, while SEQ ID NO:8

corresponds to the nucleotide sequence that is 3' to the integration site, about nucleotides 46,469 – 46,510 of the lambda phage genome.

Bacillus subtilis (BGSC #1L32, BGSC, Ohio State University, Columbus, Ohio) was chosen as an example of a gram-positive bacterial species that could be tested for the effects of a toxin-phage bacteriocide. There are many bacteriophage that are known to infect *B. subtilis*, one of which is phi-105 (BGSC #1A304(phi-105), BGSC, Ohio State University, Columbus, Ohio). The sequence of the phi-105 genome is available from the NCBI database on the Internet at ncbi.nlm.nih.gov/entrez/query.fcgi. The integration site for the TPB peptide toxin gene into the phi-105 genome was chosen to be between nucleotides 38,448 and 38,449. The nucleotide sequences of the regions immediately surrounding the chosen integration site are as follows:

GGGTAGTTGCATACCACTAAAGATGTTTCAGGTGCACATG (SEQ ID NO:9); and AGCATTGGAGGAAAGGAACGCTTTAGGGGGAAGGGAAACC (SEQ ID NO:10). SEQ ID NO:9 corresponds to the nucleotide sequence that is 5' to the integration site, about nucleotides 38,409–38,448 of the phi-105 genome, while SEQ ID NO:10 corresponds to the nucleotide sequence that is 3' to the integration site, about nucleotides 38,449–38,488 of the phi-105 genome.

Example 3: Introduction of a 3' Terminator Sequence

Before introducing the TPB peptide toxin A gene into the bacteriophage genomes, a terminator sequence can be added to the 3' end of the toxin gene in order increase the stability of toxin gene RNA synthesized within the bacterial host cell. Addition of a terminator sequence to the 3' end of the toxin gene can be accomplished by PCR, as it was in this example, as well as by other techniques known in the art, e.g., restriction fragment subcloning.

The top strand 5' oligonucleotide (SEQ ID NO:12) used to introduce the terminator sequence included the MCS sequences (SEQ ID NO:13) and a portion of the VegI/II promoter described in Example 1. The bottom strand 3' oligonucleotide (SEQ ID NO:13) used to introduce the terminator sequence included a MCS sequence distinct from the MCS sequences described in Example 1, a 3' terminator sequence, and a sequence complementary to the 3' end of the TPB peptide toxin A gene.

The top strand 5' oligonucleotide used to add the terminator sequence to the TPB peptide toxin A gene had the sequence:

CGTCCGGCGTAGAGGATCCAAGCTTTAATTAAATTTT (SEQ ID NO:11). The

underlined portion of the top strand 5' oligonucleotide sequence constitutes the MCS

sequence. The multiple cloning sites sequence was introduced to allow versatility in

manipulation of the PCR products and possible associated vectors. Restriction

endonucleases that are capable of cutting within this MCS sequence are shown in Table I.

The portion of the top strand 5' oligonucleotide sequence that is not underlined corresponds

to a portion of the VegI/II bacterial promoter added to the 5' end of the TPB peptide toxin A

gene produced in Example 1. The entire sequence of the top strand 5' oligonucleotide

sequence (SEQ ID NO:11) is capable of annealing to the TPB peptide toxin A gene

construct produced in Example 1.

The bottom strand 3' oligonucleotide used to add the terminator sequence to the TPB peptide toxin A gene had the sequence:

CGGGAAGCTTGGATCCGCATAGCAAAACGGACATCACTCCGTTTCAATGGAGGT

GATGTCCGTTTTCCGCTCGAGCTATTATTCTGGATTTCAGC (SEQ ID NO:12).

The first underlined portion of the bottom strand 3' oligonucleotide sequence constitutes the

MCS sequence. Restriction endonucleases that are capable of cutting within this MCS

sequence are shown in Table III.

Table III

Enzyme	#	Position	Sequence
AclI	1	18	ccgc
AclWI	1	15	ggatc
AluI	1	7	ag/ct
AlwI	1	15	ggatc
BamHI	1	11	g/gatcc
Bsp143I	1	11	/gatc
BstI	1	11	g/gatcc
BstX2I	1	11	r/gatcy
BstYI	1	11	r/gatcy
CviJI	1	7	rg/cy
DpnI	1	13	ga/tc
DpnII	1	11	/gatc
HindIII	1	5	a/agctt
Kzo9I	1	11	/gatc

5	MboI	1	11	/gatc
	MflI	1	11	r/gatcy
	NdeII	1	11	/gatc
	NlaIV	1	13	ggn/ncc
	PspN4I	1	13	ggn/ncc
	Sau3AI	1	11	/gatc
	XhoII	1	11	r/gatcy

The central portion of the bottom strand 3' oligonucleotide sequence (**SEQ ID NO:12**), which is not underlined above, is the 3' terminator sequence complement. The corresponding 3' terminator sequence has been shown to form a stem-loop structure that is a positive retroregulator that stabilizes mRNAs in bacteria. This 3' terminator sequence has been described in Wong and Chang (1986) *Proc Natl Acad Sci USA* 83:3233. The second underlined portion of the bottom strand 3' oligonucleotide sequence is the complement of the 3' end of the TPB peptide toxin A gene sequence (**SEQ ID NO:1**). This sequence is capable of annealing to the bottom strand of the TPB peptide toxin A gene master stock plasmid produced in Example 1, e.g., in a PCR reaction.

DNA isolated from the toxin gene bacterial stock produced in Example 1 was used as template for the PCR reaction involving the top strand 5' and bottom strand 3' oligonucleotides described above (**SEQ ID NOS:11 and 12**, respectively). Following PCR, the amplified TPB peptide toxin A gene containing the 3' terminator sequence was gel purified (Qiagen, QIAquick Gel Extraction Kit, Cat.No.28704), analyzed by endonuclease restriction fragment analysis, and used in Example 4.

Example 4: Recombinogenic Bacteriophage-Integrating Intracellular Peptide Toxin Genes

Generation of toxin genes recombinogenic with a phage genome can be produced by introducing phage genomic sequences located 5' and 3' to a chosen integration site in the phage genome to the 5' and 3' ends, respectively, of a intracellular peptide toxin encoding gene.

Generation of toxin genes recombinogenic with the lambda phage genome were produced by the addition of lambda phage genomic sequences located 5' (**SEQ ID NO:7**) and 3' (**SEQ ID NO:8**) to the chosen integration site (see Example 2) to the 5' and 3' ends, respectively, of the TPB peptide toxin A gene produced in Example 3. A single round of PCR was used to make the additions. The primers used in the PCR reaction included a 5'

Lambda Oligonucleotide (SEQ ID NO:13), consisting of a MCS sequence, a 5' homologous recombination sequence, a HindIII restriction site sequence, and a 5' annealing sequence, and a 3' Lambda Oligonucleotide (SEQ ID NO:14), consisting of a MCS sequence, a 3' homologous recombination sequence, a second MCS sequence, and a 3' annealing sequence.

5 The 5' Lambda Oligonucleotide had the sequence:

CCGGAATTCGCTAGCGGGCCCGAGTTGCCCATATCGATGGGCAACTCATGCAAT
TATTGTGAGAAGCTTTAATTAAATTTTATTGACAAAAATGGG (SEQ ID NO:13).

10 The first underlined portion of the 5' Lambda Oligonucleotide sequence constitutes MCS sequence. The MCS was introduced so that it would be easier to manipulate the PCR product for possible cloning into alternative vectors. Alternatively, the MCS region allows one to determine whether the toxin gene had integrated into the desired location in the lambda phage genome. Integration events that retain this MCS are not likely to have occurred in the desired location and can be discarded, whereas integration events that occurred via homologous recombination are likely to lack this MCS. Alternatively, the homologous recombinant sequence can be PCR amplified without the MCS and introduced into the phage genome. Restriction endonucleases that are capable of cutting within this MCS sequence are shown in Table IV.

20 Table IV

Enzyme	#	Position	Sequence
Acil	1	17	ccgc
AcsI	1	4	r/aatty
Ama87I	1	19	c/ycgrg
25 ApaI	1	20	gggcc/c
ApoI	1	4	r/aatty
AspS9I	1	16	g/gncc
AsuI	1	16	g/gncc
AvaI	1	19	c/ycgrg
30 BanII	1	20	grgcy/c
BcoI	1	19	c/ycgrg
BfaI	1	11	c/tag
BmyI	1	20	gdgch/c
BsiSI	1	1	c/cgg
35 BsoBI	1	19	c/ycgrg
Bsp120I	1	16	g/ggccc
Bsp1286I	1	20	gdgch/c

	BsuRI	1	18	gg/cc
	Cac8I	1	12	gcn/ngc
	Cfr13I	1	16	g/gncc
	CviII	1	18	rg/cy
5	Eco24I	1	20	grgcy/c
	Eco88I	1	19	c/ycgrg
	EcoRI	1	4	g/aattc
	FauI	1	18	cccgc
	FriOI	1	20	grgcy/c
10	HaeIII	1	18	gg/cc
	HapII	1	1	c/cgg
	HpaII	1	1	c/cgg
	MaeI	1	11	c/tag
	MspI	1	1	c/cgg
15	NheI	1	10	g/ctagc
	NlaIV	1	18	ggn/ncc
	Pall	1	18	gg/cc
	PspN4I	1	18	ggn/ncc
	PspOMI	1	16	g/ggccc
20	PstNHI	1	10	g/ctagc
	Sau96I	1	16	g/gncc
	SduI	1	20	gdgch/c
	Sse9I	1	4	/aatt
	Tsp509I	1	4	/aatt
25	TspEI	1	4	/aatt

The first portion of the 5' Lambda Oligonucleotide (that is not underlined) constitutes the 5' homologous recombination sequence, which was identified in Example 2 as the lambda phage sequence 5' to the integration site. The second underlined portion of the 5' Lambda Oligonucleotide constitutes a Hind III restriction site. Successful targeting of the toxin gene to the chosen site in the lambda phage genome will also result in the introduction of a new Hind III restriction site into the genome at the chosen site. Thus, restriction digest analysis of targeted lambda clones can help assess whether the targeting was successful and whether the toxin gene that has been introduced is intact, i.e., lacks deletions, rearrangements, etc. The second portion of the 5' Lambda Oligonucleotide that is not underlined constitutes the 5' annealing region, which is homologous to a portion of the VegI/II promoter sequence located at the 5' end of the PCR product produced in Example 3. This sequence is designed to anneal to the PCR product of Example 3, thereby allowing PCR amplification of a toxin gene that contains lambda phage targeting sequences.

The 3' Lambda Oligonucleotide had the sequence:

CGCCCTAGGCGGCCGAGGACCCTACTTTAGGCATTTATACTCCGCTGGAAGCGC
GTGTGTATTGGCATGCATCGATTAGTAAAACGGACATCACTCCG (SEQ ID

NO:14). The first underlined portion of the 3' Lambda Oligonucleotide sequence constitutes the first MCS sequence. The MCS was introduced so that it would be easier to manipulate the PCR product for possible cloning into alternative vectors. In addition, this multiple cloning sites sequence was introduced so that it would be easier to determine whether the toxin gene had integrated into the desired location in the lambda phage genome. Integration events that retain this MCS are not likely to have occurred in the desired location and can be discarded, whereas integration events that occurred via homologous recombination are likely to lack this MCS. Alternatively, the homologous recombinant sequence can be PCR amplified without the MCS and introduced into the phage genome. Restriction endonucleases that are capable of cutting within this MCS sequence are shown in Table V.

Table V

Enzyme	#	Position	Sequence
Acil	1	12	ccgc
AspS9I	1	17	g/gncc
AsuI	1	17	g/gncc
AvaII	1	17	g/gwcc
AvrII	1	4	c/ctagg
BfaI	1	5	c/tag
BlnI	1	4	c/ctagg
BmeI8I	1	17	g/gwcc
BsaJI	2	4,13	c/cnngg
BsaOI	1	13	cgry/cg
BseDI	2	4,13	c/cnngg
Bsh1285I	1	13	cgry/cg
BsiEI	1	13	cgry/cg
BsoFI	1	10	gc/ngc
BssTII	1	4	c/cwwgg
BstMCI	1	13	cgry/cg
BstZI	1	10	c/ggccg
BsuRI	1	12	gg/cc
Cfr13I	1	17	g/gncc
CfrI	1	10	y/ggccr
CviJI	1	12	rg/cy
DraII	1	17	rg/gnccy
EaeI	1	10	y/ggccr
EagI	1	10	c/ggccg

30

30

35

40

	BscI	1	9	at/cgat
	BseCI	1	9	at/cgat
	Bsp106I	1	9	at/cgat
	BspDI	1	9	at/cgat
5	BspXI	1	9	at/cgat
	Bsu15I	1	9	at/cgat
	Cac8I	1	4	gcn/ngc
	ClaI	1	9	at/cgat
	EcoT22I	1	8	atgca/t
10	Hsp92II	1	6	catg/
	Mph1103I	1	8	atgca/t
	NlaIII	1	6	catg/
	NsiI	1	8	atgca/t
	NspI	1	6	rcatg/y
15	PaeI	1	6	gcatg/c
	Ppu10I	1	4	a/tgcat
	SfaNI	1	10	gcatc
	SphI	1	6	gcatg/c
	TaqI	1	9	t/cga
20	TthHB8I	1	9	t/cga
	Zsp2I	1	8	atgca/t

The second portion of the 3' Lambda Oligonucleotide (SEQ ID NO:25) that is not underlined constitutes the 5' annealing region (SEQ ID NO:28), which is complementary to a portion of the 3' terminator sequence located at the 3' end of the PCR product produced in Example 3. This sequence is designed to anneal to the PCR product of Example 3, thereby allowing PCR amplification of a TPB peptide toxin A gene that contains lambda phage targeting sequences.

Generation of toxin genes recombinagenic with the phi-105 phage genome were produced by the addition of phi-105 phage genomic sequences located 5'(SEQ ID NO:9) and 3' (SEQ ID NO:10) to the chosen integration site (see Example 2) to the 5' and 3' ends, respectively, of the TPB peptide toxin A gene produced in Example 3. A single round of PCR was used to make the additions. The primers used in the PCR reaction included a 5' Phi-105 Oligonucleotide (SEQ ID NO:15), consisting of a MCS sequence, a 5' homologous recombination sequence, a HindIII restriction site sequence, and a 5' annealing sequence, and a 3' Phi-105 Oligonucleotide (SEQ ID NO:16), consisting of a MCS sequence, a 3' homologous recombination sequence, a second MCS sequence, and a 3' annealing sequence.

The 5' Phi-105 Oligonucleotide had the sequence:

CCGGAATTCGCTAGCGGGCCCGAGGGGTAGTTGCATACCACTAAAGATGTTTCAG
GTGCACATGAAGCTTTAATTAAATTTTATTTGACAAAAATGGG (SEQ ID NO:15).

The first underlined portion of the 5' Phi-105 Oligonucleotide sequence constitutes the MCS sequence. The multiple cloning sites sequence was introduced so that it would be easier to
 5 manipulate the PCR product for possible cloning into alternative vectors. In addition, the MCS region allows one to determine whether the toxin gene had integrated into the desired location in the lambda phage genome. Integration events that retain this MCS are not likely to have occurred in the desired location and can be discarded, whereas integration events that occurred via homologous recombination are likely to lack this MCS. Alternatively, the
 10 homologous recombinant sequence can be PCR amplified without the MCS and introduced into the phage genome. Restriction endonucleases that are capable of cutting within this MCS sequence are shown in Table VII.

Table VII

	Enzyme	#	Position	Sequence
5	Acil	1	17	ccgc
	AcsI	1	4	r/aatty
	Ama87I	1	19	c/ycgrg
	ApaI	1	20	gggcc/c
10	ApoI	1	4	r/aatty
	AspS9I	1	16	g/gncc
	AsuI	1	16	g/gncc
	AvaI	1	19	c/ycgrg
	BanII	1	20	grgcy/c
	BcoI	1	19	c/ycgrg
	BfaI	1	11	c/tag
	BmyI	1	20	gdgch/c
	BsiSI	1	1	c/cgg
	BsoBI	1	19	c/ycgrg
30	Bsp120I	1	16	g/ggccc
	Bsp1286I	1	20	gdgch/c
	BsuRI	1	18	gg/cc
	Cac8I	1	12	gcn/ngc
	Cfr13I	1	16	g/gncc
35	CviJI	1	18	rg/cy
	Eco24I	1	20	grgcy/c
	Eco88I	1	19	c/ycgrg
	EcoRI	1	4	g/aattc
	FauI	1	18	cccgc
40	FriOI	1	20	grgcy/c

	HaeIII	1	18	gg/cc
	HapII	1	1	c/cgg
	HpaII	1	1	c/cgg
	MaeI	1	11	c/tag
5	MspI	1	1	c/cgg
	NheI	1	10	g/ctagc
	NlaIV	1	18	ggn/ncc
	Pall	1	18	gg/cc
	PspN4I	1	18	ggn/ncc
10	PspOMI	1	16	g/ggccc
	PstNHI	1	10	g/ctagc
	Sau96I	1	16	g/gncc
	SduI	1	20	gdgch/c
	Sse9I	1	4	/aatt
15	Tsp509I	1	4	/aatt
	TspEI	1	4	/aatt

The first portion of the 5' Phi-105 Oligonucleotide that is not underlined constitutes the 5' homologous recombination sequence, which was identified in Example 2 as the phi-105 phage sequence 5' to the integration site. The second underlined portion of the 5' Phi-105 Oligonucleotide constitutes a Hind III restriction site. Successful targeting of the toxin gene to the chosen site in the phi-105 phage genome will also result in the introduction of a new Hind III restriction site into the genome at the chosen site. Thus, restriction digest analysis of targeted phi-105 clones can help assess whether the targeting was successful and whether the toxin gene that has been introduced is intact, i.e., lacks deletions, rearrangements, etc. The second portion of the 5' Phi-105 Oligonucleotide that is not underlined constitutes the 5' annealing region, which is homologous to a portion of the VegI/II promoter sequence located at the 5' end of the PCR product produced in Example 3. This sequence is designed to anneal to the PCR product of Example 3, thereby allowing PCR amplification of a toxin gene that contains phi-105 phage targeting sequences.

The 3' Phi-105 Oligonucleotide had the following sequence:

CGCCCTAGGCGGCCGAGGACCCGGTTTCCCTTCCCCCTAAAGCGTTCCTTTCCTC
CAATGCTGGCATGCATCGATTAGTAAAACGGACATCACTCCG (SEQ ID NO:16).

The first underlined portion of the 3' Phi-105 Oligonucleotide sequence constitutes the first MCS sequence. This multiple cloning site sequence was introduced so that it would be easier to manipulate the PCR product for possible cloning into alternative vectors. In addition, it would be easier to determine whether the toxin gene had integrated into the desired location

in the lambda phage genome. Integration events that retain this MCS are not likely to have occurred in the desired location and can be discarded, whereas integration events that occurred via homologous recombination are likely to lack this MCS. Alternatively, the homologous recombinant sequence can be PCR amplified without the MCS and introduced into the phage genome. Restriction endonucleases that are capable of cutting within this MCS sequence are shown in Table VIII.

Table VIII

Enzyme	#	Position	Sequence
AciI	1	12	ccgc
AspS9I	1	17	g/gncc
AsuI	1	17	g/gncc
AvaII	1	17	g/gwcc
AvrII	1	4	c/ctagg
BfaI	1	5	c/tag
BlnI	1	4	c/ctagg
Bme18I	1	17	g/gwcc
BsaJI	2	4, 13	c/cnngg
BsaOI	1	13	cgry/cg
BseDI	2	4, 13	c/cnngg
Bsh1285I	1	13	cgry/cg
BsiEI	1	13	cgry/cg
BsoFI	1	10	gc/ngc
BssTII	1	4	c/cwwgg
BstMCI	1	13	cgry/cg
BstZI	1	10	c/ggccg
BsuRI	1	12	gg/cc
Cfr13I	1	17	g/gncc
CfrI	1	10	y/ggcer
CviJI	1	12	rg/cy
DraII	1	17	rg/gnccy
EaeI	1	10	y/ggcer
EagI	1	10	c/ggccg
EclXI	1	10	c/ggccg
Eco130I	1	4	c/cwwgg
Eco47I	1	17	g/gwcc
Eco52I	1	10	c/ggccg
EcoO109I	1	17	rg/gnccy
EcoT14	1	4	c/cwwgg
ErhI	1	4	c/cwwgg
Fsp4HI	1	10	gc/ngc
HaeIII	1	12	gg/cc

	HgiEI	1	17	g/gwcc
	ItaI	1	10	gc/ngc
	MaeI	1	5	c/tag
	MnII	1	18	cctc
5	NlaIV	1	19	ggn/ncc
	PalI	1	12	gg/cc
	PpuMI	1	17	rg/gwccy
	Psp5II	1	17	rg/gwccy
	PspN4I	1	19	ggn/ncc
10	Sau96I	1	17	g/gncc
	SinI	1	17	g/gwcc
	StyI	1	4	c/cwwgg
	XmaIII	1	10	c/ggccg

15 The first portion of the 3' Phi-105 Oligonucleotide that is not underlined constitutes the 3' homologous recombination sequence, which was identified in Example 2 as the phi-105 phage sequence 3' to the integration site. The second underlined portion of the 3' Phi-105 Oligonucleotide constitutes a second MCS sequence. Successful targeting of the toxin gene to the chosen site in the phi-105 phage genome will also result in the introduction of the restriction sites present in this MCS into the genome at the chosen site. Thus, restriction digest analysis of targeted phi-105 clones can help assess whether the targeting was successful and whether the toxin gene that has been introduced is intact, i.e., lacks deletions, rearrangements, etc. Restriction endonucleases that are capable of cutting within this MCS sequence are shown in Table IX.

Table IX

	Enzyme	#	Position	Sequence
	BanIII	1	9	at/cgat
	BbuI	1	6	gcatg/c
30	Bsa29I	1	9	at/cgat
	BscI	1	9	at/cgat
	BseCI	1	9	at/cgat
	Bsp106I	1	9	at/cgat
	BspDI	1	9	at/cgat
35	BspXI	1	9	at/cgat
	Bsu15I	1	9	at/cgat
	Cac8I	1	4	gcn/ngc
	ClaI	1	9	at/cgat
	EcoT22I	1	8	atgca/t
40	Hsp92II	1	6	catg/

	Mph1103I	1	8	atgca/t
	NlaIII	1	6	catg/
	NsiI	1	8	atgca/t
	NspI	1	6	rcatg/y
5	PaeI	1	6	gcatg/c
	Ppu10I	1	4	a/tgcat
	SfaNI	1	10	gcatc
	SphI	1	6	gcatg/c
	TaqI	1	9	t/cga
10	TthHB8I	1	9	t/cga
	Zsp2I	1	8	atgca/t

The second portion of the 3' Phi-105 Oligonucleotide that is not underlined constitutes the 5' annealing region, which is complementary to a portion of the 3' terminator sequence located at the 3' end of the PCR product produced in Example 3. This sequence is designed to anneal to the PCR product of Example 3, thereby allowing PCR amplification of a TPB peptide toxin A gene that contains phi-105 phage targeting sequences.

Example 5: *In vitro* homologous recombination of an intracellular peptide toxin gene into a bacteriophage genome

The homologous recombination event can be manipulated *in vitro* using isolated bacteriophage DNA added to a bacterial host cell supernatant seeded with the homologous recombination competent PCR product containing a intracellular toxin gene, using the basic protocols as described by Mackal, et al (1964), *PNAS* 51:1172, the contents of which are incorporated herein by reference. Such a procedure was performed with the recombinogenic TPB peptide toxin gene PCR products produced in Example 4. After incubation at 37°C, the reaction mixtures are added to cell cultures of host cells and plated on the appropriate media by mixing in 2 mls of molten top agar poured onto a hardened bottom agar. Plates are incubated at 37° for the *E coli* C600 gram-negative bacteria, and 30°C for the *B. subtilis* 1L32 gram-positive bacteria. Plaques are screened for incorporation of the TPB peptide using Southern hybridization techniques. Plaques identified as positive are isolated and stocks are prepared from the single plaques. Chromosomal DNA isolated from these stocks is analyzed by restriction digestion, followed by sequencing.

Example 6: *In vivo* homologous recombination of an intracellular peptide toxin gene into a bacteriophage genome

The homologous recombination event can be manipulated *in vivo* with competent bacterial cells lysogenic for the chosen bacteriophage. Competent *E coli* C600 gram-negative bacteria lysogenic for the wild-type lambda phage are prepared using the calcium chloride method, as described in Molecular Cloning (1989), 2nd Ed., Sambrook et al., Eds., Cold Spring Harbor Press, the contents of which are incorporated herein by reference. Competent *B. Subtilis* 1L32 lysogenic for phi-105 phage are prepared using methods described by Errington & Mandelstam (1983), *Journal of General Microbiology* 129:2091, the contents of which are incorporated herein by reference. The recombinagenic PCR product containing the intracellular peptide toxin gene is added to the competent bacterial cells and heat shocked as described in Sambrook et al., supra. After a one hour incubation at 37°C, the reaction mixtures are added to cell cultures of host cells and plated on the appropriate media by mixing in 2 mls of molten top agar poured onto a hardened bottom agar. Plates are incubated at 37° for the *E coli* C600 gram-negative bacteria, and 30°C for the *B. subtilis* 1L32 gram-positive bacteria. Plaques are screened for incorporation of the TPB peptide using Southern hybridization techniques. Plaques identified as positive are isolated and stocks are prepared from the single plaques. Chromosomal DNA isolated from these stocks is analyzed by restriction digestion, followed by sequencing.

Example 7: Use of a Toxin-Phage Bacteriocide to kill bacteria

To test the effectiveness of TPB peptide toxin A, lambda phage (American Type Culture Collection (ATCC) Accession No. 23724-B2) was engineered to express TPB peptide toxin A. This modified phage killed 100% of *E. coli* (ATCC Accession No. 23724). No lysogenic colonies were observed.

In addition, phi-105 phage (BGSC Accession No. 1A304 (phi105); Ohio State University, Columbus, OH) was engineered to express TPB peptide toxin A. This modified phage killed 100% of *B. subtilis* (BGSC Accession No. 1L32). No lysogenic clones were observed.

A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention. Accordingly, other embodiments are within the scope of the

5 following claims.